arXiv:1403.0435v1 [math.NT] 3 Mar 2014

RECURRENCE FORMULAS WITH GAPS FOR BERNOULLI AND EULER POLYNOMIALS

ZHI-HONG SUN

School of Mathematical Sciences, Huaiyin Normal University,
Huaian, Jiangsu 223001, PR China
E-mail: zhihongsun@yahoo.com
Homepage: http://www.hytc.edu.cn/xsjl/szh

ABSTRACT. Let $\{B_n\}$, $\{B_n(x)\}$ and $\{E_n(x)\}$ be the Bernoulli numbers, Bernoulli polynomials and Euler polynomials, respectively. In this paper we mainly establish formulas for $\sum_{6|k-3}\binom{n}{k}B_{n-k}(x)$, $\sum_{6|k}\binom{n}{k}E_{n-k}(x)$ and $\sum_{6|k-3}\binom{n}{k}m^kB_{n-k}$ in the cases $m=2,3,4$.

MSC: Primary 11B68, Secondary 11B39
Keywords: Bernoulli number; Bernoulli polynomial; Euler polynomial

1. Introduction.

The Bernoulli numbers $\{B_n\}$ and Bernoulli polynomials $\{B_n(x)\}$ are defined by

$$B_0=1,\ \sum_{k=0}^{n-1}\binom{n}{k}B_k=0\ (n\ge 2)\quad\text{and}\quad B_n(x)=\sum_{k=0}^{n}\binom{n}{k}B_kx^{n-k}\ (n\ge 0).$$

The Euler numbers $\{E_n\}$ and Euler polynomials $\{E_n(x)\}$ are defined by

$$\frac{2\mathrm{e}^t}{\mathrm{e}^{2t}+1}=\sum_{n=0}^{\infty}E_n\frac{t^n}{n!}\ (|t|<\frac{\pi}{2})\quad\text{and}\quad\frac{2\mathrm{e}^{xt}}{\mathrm{e}^t+1}=\sum_{n=0}^{\infty}E_n(x)\frac{t^n}{n!}\ (|t|<\pi),\tag{1.1}$$

which are equivalent to (see [MOS])

$$E_0=1,\ E_{2n-1}=0,\ \sum_{r=0}^{n}\binom{2n}{2r}E_{2r}=0\ (n\ge 1)$$

and

$$E_n(x)+\sum_{r=0}^{n}\binom{n}{r}E_r(x)=2x^n\ (n\ge 0).\tag{1.2}$$

The first few Bernoulli and Euler numbers are shown below:

$$B_0=1,\ B_1=-\frac{1}{2},\ B_2=\frac{1}{6},\ B_4=-\frac{1}{30},\ B_6=\frac{1}{42},\ B_8=-\frac{1}{30},B_{10}=\frac{5}{66},$$
$$E_0=1,\ E_2=-1,\ E_4=5,\ E_6=-61,\ E_8=1385,\ E_{10}=-50521.$$

It is well known that ([MOS])

$$\begin{aligned} E_n(x) &= \frac{1}{2^n}\sum_{r=0}^{n}\binom{n}{r}(2x-1)^{n-r}E_r \\ &= \frac{2}{n+1}\Big(B_{n+1}(x) - 2^{n+1}B_{n+1}\Big(\frac{x}{2}\Big)\Big) \\ &= \frac{2^{n+1}}{n+1}\Big(B_{n+1}\Big(\frac{x+1}{2}\Big) - B_{n+1}\Big(\frac{x}{2}\Big)\Big). \end{aligned} \tag{1.3}$$

In particular,

$$E_n = 2^n E_n\Big(\frac{1}{2}\Big) \quad \text{and} \quad E_n(0) = \frac{2(1-2^{n+1})B_{n+1}}{n+1}. \tag{1.4}$$

In his first paper Ramanujan [R] found some recurrence formulas with gaps for Bernoulli numbers. In particular, he showed that for $n = 3, 5, 7, \dots$,

$$\sum_{\substack{k=0\\6|k-3}}^{n}\binom{n}{k}B_{n-k} = \begin{cases} -\frac{n}{6} & \text{if } n \equiv 1 \pmod 6, \\ \frac{n}{3} & \text{if } n \equiv 3, 5 \pmod 6. \end{cases} \tag{1.5}$$

Some generalizations of (1.5) were given by M. Chellali [C] in 1988. However, the proofs of (1.5) given by Ramanujan and Chellali are somewhat complicated. Let $\omega = (-1+\sqrt{-3})/2$. In this paper, we prove the following generalization of (1.5) for Bernoulli polynomials in a very simple manner:

$$\sum_{\substack{k=0\\6|k-3}}^{n}\binom{n}{k}B_{n-k}(x) = \frac{n}{6}\big(x^{n-1} + (x-1)^{n-1} - (x+\omega)^{n-1} - (x+\omega^2)^{n-1}\big).$$

As consequences we give explicit formulas for $\sum\limits_{\substack{k=0\\6|k-3}}^{n}\binom{n}{k}m^k B_{n-k}$ in the cases $m = 2, 3, 4$.

Let $\mathbb{N}$ be the set of positive integers, and let $[x]$ be the greatest integer not exceeding x. In [L] Lehmer showed that for $n = 2, 4, 6, \dots$,

$$4E_n = 2\big(1 + (-3)^{\frac{n}{2}}\big) - 3\sum_{k=1}^{[n/6]}\binom{n}{6k}2^{6k}E_{n-6k}. \tag{1.6}$$

In this paper we prove that for $n \in \mathbb{N}$,

$$4E_n(x) + 3\sum_{k=1}^{[n/6]}\binom{n}{6k}E_{n-6k}(x) = x^n + (x-1)^n + (x+\omega)^n + (x+\omega^2)^n.$$

2. Recurrence formulas with gaps for Bernoulli polynomials.

For two numbers b and c, let $\{U_n(b,c)\}$ and $\{V_n(b,c)\}$ be the Lucas sequences given by

$$U_0(b,c)=0,\ U_1(b,c)=1,\ U_{n+1}(b,c)=bU_n(b,c)-cU_{n-1}(b,c)\ (n\geq 1)$$

and

$$V_0(b,c)=2,\ V_1(b,c)=b,\ V_{n+1}(b,c)=bV_n(b,c)-cV_{n-1}(b,c)\ (n\geq 1).$$

It is well known that (see [W]) for $b^2-4c\neq 0$,

$$U_n(b,c)=\frac{1}{\sqrt{b^2-4c}}\Big\{\Big(\frac{b+\sqrt{b^2-4c}}{2}\Big)^n-\Big(\frac{b-\sqrt{b^2-4c}}{2}\Big)^n\Big\}$$

and

$$V_n(b,c)=\Big(\frac{b+\sqrt{b^2-4c}}{2}\Big)^n+\Big(\frac{b-\sqrt{b^2-4c}}{2}\Big)^n.$$

Lemma 2.1. *For $n\in\mathbb{N}$ we have*

$$\sum_{k=0}^{n-1}\binom{n}{k}B_k(x)((1+z)^{n-k}-z^{n-k})=n(x+z)^{n-1}.$$

Proof. It is known that (see [MOS])

$$\sum_{k=0}^{n}\binom{n}{k}B_k(x)y^{n-k}=B_n(x+y)\quad\text{and}\quad B_n(t+1)-B_n(t)=nt^{n-1}. \tag{2.1}$$

Thus,

$$\begin{aligned}&\sum_{k=0}^{n-1}\binom{n}{k}B_k(x)((1+z)^{n-k}-z^{n-k})\\&=\sum_{k=0}^{n}\binom{n}{k}B_k(x)(1+z)^{n-k}-B_n(x)-\sum_{k=0}^{n}\binom{n}{k}B_k(x)z^{n-k}+B_n(x)\\&=B_n(x+z+1)-B_n(x+z)=n(x+z)^{n-1}.\end{aligned}$$

This proves the lemma.

Theorem 2.1. *For $n\in\mathbb{N}$ we have*

$$\sum_{\substack{k=0\\6|k-3}}^{n}\binom{n}{k}B_{n-k}(x)=\frac{n}{6}\big(x^{n-1}+(x-1)^{n-1}-(x+\omega)^{n-1}-(x+\omega^2)^{n-1}\big).$$

Proof. As $1+\omega+\omega^2=0$ and $\omega^3=1$, using Lemma 2.1 we see that

$$\begin{aligned}
&-4\sum_{\substack{k=0\\ n-k\equiv 3 \pmod 6}}^{n}\binom{n}{k}B_k(x)+2\sum_{\substack{k=0\\ n-k\equiv \pm 1 \pmod 6}}^{n}\binom{n}{k}B_k(x)\\
&=\sum_{k=0}^{n-1}\binom{n}{k}B_k(x)((-1)^{n-k}-1)(\omega^{n-k}+\omega^{2(n-k)})\\
&=\sum_{k=0}^{n-1}\binom{n}{k}B_k(x)((-\omega^2)^{n-k}-\omega^{n-k}+(-\omega)^{n-k}-\omega^{2(n-k)})\\
&=\sum_{k=0}^{n-1}\binom{n}{k}B_k(x)((1+\omega)^{n-k}-\omega^{n-k}+(1+\omega^2)^{n-k}-\omega^{2(n-k)}))\\
&=n(x+\omega)^{n-1}+n(x+\omega^2)^{n-1}.
\end{aligned}$$

That is,

$$-6\sum_{\substack{k=0\\ 6|n-k-3}}^{n}\binom{n}{k}B_k(x)+2\sum_{\substack{k=0\\ 2\nmid n-k}}^{n}\binom{n}{k}B_k(x)=n(x+\omega)^{n-1}+n(x+\omega^2)^{n-1}.$$

From Lemma 2.1 we have

$$\text{(2.2)}\quad \sum_{k=0}^{n-1}\binom{n}{k}B_k(x)=nx^{n-1} \text{ and } \sum_{k=0}^{n-1}\binom{n}{k}B_k(x)(-1)^{n-k+1}=n(x-1)^{n-1}.$$

Thus,

$$2\sum_{\substack{k=0\\ 2\nmid n-k}}^{n}\binom{n}{k}B_k(x)=\sum_{k=0}^{n-1}\binom{n}{k}B_k(x)(1-(-1)^{n-k})=nx^{n-1}+n(x-1)^{n-1}.$$

Hence

$$\begin{aligned}
6\sum_{\substack{k=0\\ 6|k-3}}^{n}\binom{n}{k}B_{n-k}(x)&=6\sum_{\substack{r=0\\ 6|n-r-3}}^{n}\binom{n}{r}B_r(x)\\
&=nx^{n-1}+n(x-1)^{n-1}-n(x+\omega)^{n-1}-n(x+\omega^2)^{n-1}.
\end{aligned}$$

This completes the proof.

As $B_n(0)=B_n$, taking $x=0$ in Theorem 2.1 we obtain Ramanujan's identity (1.5).

Theorem 2.2. *For $m \in \mathbb{N}$ and $n = 3, 5, 7, \ldots$ we have*

$$\sum_{\substack{k=0\\6|k-3}}^{n} \binom{n}{k} m^k B_{n-k}$$
$$= \frac{n}{3}\Big\{ \sum_{r=1}^{m-1} r^{n-1} - \sum_{r=1}^{[\frac{m-1}{2}]} \big((r+m\omega)^{n-1} + (r+m\omega^2)^{n-1}\big) + \delta(m,n)\Big\},$$

where

$$\delta(m,n) = \begin{cases} -\frac{1}{2}m^{n-1} & \text{if } 2 \nmid m \text{ and } 3 \mid n-1, \\ m^{n-1} & \text{if } 2 \nmid m \text{ and } 3 \nmid n-1, \\ -\frac{1}{2}m^{n-1} - (-3)^{\frac{n-1}{2}}(\frac{m}{2})^{n-1} & \text{if } 2 \mid m \text{ and } 3 \mid n-1, \\ m^{n-1} - (-3)^{\frac{n-1}{2}}(\frac{m}{2})^{n-1} & \text{if } 2 \mid m \text{ and } 3 \nmid n-1. \end{cases}$$

Proof. From Theorem 2.1 we have

$$\sum_{\substack{k=0\\6|k-3}}^{n} \binom{n}{k} \sum_{r=0}^{m-1} B_{n-k}\Big(\frac{r}{m}\Big)$$
$$= \frac{n}{6}\sum_{r=0}^{m-1} \Big\{\Big(\frac{r}{m}\Big)^{n-1} + \Big(\frac{r-m}{m}\Big)^{n-1} - \Big(\frac{r+m\omega}{m}\Big)^{n-1} - \Big(\frac{r+m\omega^2}{m}\Big)^{n-1}\Big\}.$$

By Raabe's theorem (see [MOS]), $\sum_{r=0}^{m-1} B_{n-k}(\frac{r}{m}) = m^{1-(n-k)}B_{n-k}$. Thus,

$$\sum_{\substack{k=0\\6|k-3}}^{n} \binom{n}{k} m^k B_{n-k}$$
$$= \frac{n}{6}\sum_{r=0}^{m-1} \Big\{r^{n-1} + (m-r)^{n-1} - (r+m\omega)^{n-1} - (r+m\omega^2)^{n-1}\Big\}$$
$$= \frac{n}{6}\Big\{2\sum_{r=1}^{m-1} r^{n-1} + m^{n-1} - m^{n-1}(\omega^{n-1} + \omega^{2(n-1)})$$
$$- \sum_{r=1}^{m-1}((r+m\omega)^{n-1} + (r+m\omega^2)^{n-1})\Big\}.$$

As $2 \mid n-1$ and $1 + \omega + \omega^2 = 0$, we see that

$$(m-r+m\omega)^{n-1} + (m-r+m\omega^2)^{n-1} = (r+m\omega)^{n-1} + (r+m\omega^2)^{n-1}.$$

Hence

$$\sum_{\substack{k=0\\6|k-3}}^{n}\binom{n}{k}m^kB_{n-k}$$
$$=\frac{n}{6}\Big\{m^{n-1}(1-\omega^{n-1}-\omega^{2(n-1)})-(1+(-1)^m)\Big(\frac{m}{2}\Big)^{n-1}(-3)^{\frac{n-1}{2}}$$
$$+2\sum_{r=1}^{m-1}r^{n-1}-2\sum_{r=1}^{[(m-1)/2]}\Big((r+m\omega)^{n-1}+(r+m\omega^2)^{n-1}\Big)\Big\}.$$

This yields the result.

Putting $m=2$ in Theorem 2.2 we deduce the following result.

Corollary 2.1. *For $n=3,5,7,\ldots$ we have*

$$\sum_{\substack{k=0\\6|k-3}}^{n}\binom{n}{k}2^kB_{n-k}=\begin{cases}\frac{n}{3}(1-2^{n-2}-(-3)^{\frac{n-1}{2}}) & \text{if } n\equiv 1 \pmod 6,\\ \frac{n}{3}(1+2^{n-1}-(-3)^{\frac{n-1}{2}}) & \text{if } n\equiv 3,5 \pmod 6.\end{cases}$$

Corollary 2.2. *For $n=3,5,7,\ldots$ we have*

$$\sum_{\substack{k=0\\6|k-3}}^{n}\binom{n}{k}3^kB_{n-k}=\begin{cases}\frac{n}{3}\big(1+2^{n-1}-\frac{3^{n-1}}{2}-V_{n-1}(1,7)\big) & \text{if } 6\mid n-1,\\ \frac{n}{3}\big(1+2^{n-1}+3^{n-1}-V_{n-1}(1,7)\big) & \text{if } 6\nmid n-1.\end{cases}$$

Proof. Observe that

$$\begin{aligned}V_{n-1}(1,7)&=\Big(\frac{1+\sqrt{1^2-4\cdot 7}}{2}\Big)^{n-1}+\Big(\frac{1-\sqrt{1^2-4\cdot 7}}{2}\Big)^{n-1}\\&=\Big(\frac{1+3\sqrt{-3}}{2}\Big)^{n-1}+\Big(\frac{1-3\sqrt{-3}}{2}\Big)^{n-1}\\&=(1+3\omega)^{n-1}+(1+3\omega^2)^{n-1}.\end{aligned}$$

Taking $m=3$ in Theorem 2.2 we deduce the result.

Corollary 2.3. *For $n=3,5,7,\ldots$ we have*

$$\sum_{\substack{k=0\\6|k-3}}^{n}\binom{n}{k}4^kB_{n-k}$$
$$=\begin{cases}\frac{n}{3}\big(1+2^{n-1}+3^{n-1}-2^{2n-3}-(-12)^{\frac{n-1}{2}}-V_{n-1}(2,13)\big) & \text{if } 6\mid n-1,\\ \frac{n}{3}\big(1+2^{n-1}+3^{n-1}+4^{n-1}-(-12)^{\frac{n-1}{2}}-V_{n-1}(2,13)\big) & \text{if } 6\nmid n-1.\end{cases}$$

Proof. Note that

$$\begin{aligned}V_{n-1}(2,13)&=\Big(\frac{2+\sqrt{2^2-4\cdot 13}}{2}\Big)^{n-1}+\Big(\frac{2-\sqrt{2^2-4\cdot 13}}{2}\Big)^{n-1}\\&=(1+2\sqrt{-3})^{n-1}+(1-2\sqrt{-3})^{n-1}\\&=(1+4\omega)^{n-1}+(1+4\omega^2)^{n-1}.\end{aligned}$$

Taking $m=4$ in Theorem 2.2 we deduce the result.

Theorem 2.3. *For $n \in \mathbb{N}$ we have*

$$\sum_{\substack{k=0\\4|k-2}}^{n} \binom{n}{k} B_{n-k}(x)(-1)^{\frac{k-2}{4}} 2^{n+1-\frac{k}{2}} = ni\big((2x-1-i)^{n-1} - (2x-1+i)^{n-1}\big).$$

Proof. Using Lemma 2.1 we see that

$$\begin{aligned}
&4 \sum_{\substack{k=0\\4|n-k-2}}^{n-1} \binom{n}{k} B_k(x) \Big(\frac{i}{2}\Big)^{\frac{n-k}{2}} \\
&= \sum_{\substack{k=0\\2|n-k}}^{n-1} \binom{n}{k} B_k(x) \cdot 2\Big(\Big(\frac{i}{2}\Big)^{\frac{n-k}{2}} - \Big(-\frac{i}{2}\Big)^{\frac{n-k}{2}}\Big) \\
&= \sum_{k=0}^{n-1} \binom{n}{k} B_k(x)(1+(-1)^{n-k})\Big(\Big(\frac{1+i}{2}\Big)^{n-k} - \Big(\frac{i-1}{2}\Big)^{n-k}\Big) \\
&= \sum_{k=0}^{n-1} \binom{n}{k} B_k(x)\Big(\Big(\frac{1+i}{2}\Big)^{n-k} - \Big(\frac{i-1}{2}\Big)^{n-k} - \Big(\frac{1-i}{2}\Big)^{n-k} + \Big(\frac{-i-1}{2}\Big)^{n-k}\Big) \\
&= n\Big(x+\frac{i-1}{2}\Big)^{n-1} - n\Big(x+\frac{-i-1}{2}\Big)^{n-1}.
\end{aligned}$$

Thus,

$$\begin{aligned}
&\sum_{\substack{k=0\\4|k-2}}^{n} \binom{n}{k} B_{n-k}(x)(-1)^{\frac{k-2}{4}} 2^{-\frac{k-2}{2}} \\
&= \sum_{\substack{k=0\\4|k-2}}^{n} \binom{n}{k} B_{n-k}(x)\Big(\frac{i}{2}\Big)^{\frac{k-2}{2}} = \sum_{\substack{r=0\\4|n-r-2}}^{n} \binom{n}{r} B_r(x)\Big(\frac{i}{2}\Big)^{\frac{n-r}{2}-1} \\
&= \frac{n}{2i} \cdot \frac{(2x-1+i)^{n-1} - (2x-1-i)^{n-1}}{2^{n-1}}.
\end{aligned}$$

This yields the result.

Theorem 2.4. *Let $m \in \mathbb{N}$ and $n \in \{2,4,6,\dots\}$. Then*

$$\begin{aligned}
&\sum_{\substack{k=0\\4|k-2}}^{n} \binom{n}{k} (-1)^{\frac{k-2}{4}} 2^{n-\frac{k}{2}} m^k B_{n-k} \\
&= \frac{n}{2}\Big\{ m^{n-1}\Big((-1)^{[\frac{n-2}{4}]} 2^{\frac{n}{2}} - (1+(-1)^m)(-1)^{\frac{n}{2}}\Big) \\
&\quad + 2i \sum_{r=1}^{[\frac{m-1}{2}]} \Big((2r-m-mi)^{n-1} - (2r-m+mi)^{n-1}\Big)\Big\}.
\end{aligned}$$

Proof. By Raabe's theorem, $\sum_{r=0}^{m-1} B_{n-k}\big(\frac{r}{m}\big) = m^{k-(n-1)} B_{n-k}$. Thus, using Theorem 2.3 we see that

$$\begin{aligned}
&\sum_{\substack{k=0\\4|k-2}}^{n} \binom{n}{k} (-1)^{\frac{k-2}{4}} 2^{n-\frac{k}{2}} m^k B_{n-k} \\
&= \sum_{\substack{k=0\\4|k-2}}^{n} \binom{n}{k} (-1)^{\frac{k-2}{4}} 2^{n-\frac{k}{2}} m^{n-1} \sum_{r=0}^{m-1} B_{n-k}\Big(\frac{r}{m}\Big) \\
&= \frac{n}{2} i \sum_{r=0}^{m-1} \Big\{ (2r-m-mi)^{n-1} - (2r-m+mi)^{n-1} \Big\} \\
&= \frac{n}{2} i \Big\{ (-m-mi)^{n-1} - (-m+mi)^{n-1} \\
&\quad + \sum_{r=1}^{m-1} \big((2r-m-mi)^{n-1} - (2r-m+mi)^{n-1} \big) \Big\} \\
&= \frac{n}{2} i \Big\{ (-m-mi)^{n-1} - (-m+mi)^{n-1} - \big(1+(-1)^m\big)(mi)^{n-1} \\
&\quad + 2 \sum_{r=1}^{[\frac{m-1}{2}]} \big((2r-m-mi)^{n-1} - (2r-m+mi)^{n-1} \big) \Big\}
\end{aligned}$$

It is evident that

$$\begin{aligned}
&i\big((-1-i)^{n-1} - (-1+i)^{n-1} \big) \\
&= i\big((-1-i)(2i)^{\frac{n-2}{2}} - (-1+i)(-2i)^{\frac{n-2}{2}} \big) \\
&= \begin{cases} i \cdot (-2i)(2i)^{\frac{n-2}{2}} = (-1)^{\frac{n-2}{4}} 2^{\frac{n}{2}} & \text{if } 4 \mid n-2, \\ i \cdot (-2)(2i)^{\frac{n-2}{2}} = (-1)^{[\frac{n-2}{4}]} 2^{\frac{n}{2}} & \text{if } 4 \mid n \end{cases}
\end{aligned}$$

and

$$\begin{aligned}
&(2(m-r)-m-mi)^{n-1} - (2(m-r)-m+mi)^{n-1} \\
&= (2r-m-mi)^{n-1} - (2r-m+mi)^{n-1}.
\end{aligned}$$

Now combining all the above we deduce the result.

Putting $m = 1, 2$ in Theorem 2.4 we deduce the following identities.

Corollary 2.4. *(Ramanujan [R]) For $n = 2, 4, 6, \ldots$ we have*

$$\sum_{\substack{k=0\\4|k-2}}^{n} \binom{n}{k} (-1)^{\frac{k-2}{4}} 2^{\frac{n-k}{2}} B_{n-k} = (-1)^{[\frac{n-2}{4}]} \frac{n}{2}.$$

Corollary 2.5. *For $n = 2, 4, 6, \ldots$ we have*

$$\sum_{\substack{k=0\\4|k-2}}^{n} \binom{n}{k} (-1)^{\frac{k-2}{4}} 2^{\frac{k}{2}} B_{n-k} = \frac{n}{2}\big((-1)^{[\frac{n-2}{4}]} 2^{\frac{n-2}{2}} + (-1)^{\frac{n-2}{2}}\big).$$

Corollary 2.6. *For $n = 2, 4, 6, \ldots$ we have*

$$\sum_{\substack{k=0\\4|k-2}}^{n} \binom{n}{k} (-1)^{\frac{k-2}{4}} 2^{\frac{3k}{2}} B_{n-k} = \frac{n}{2}\big((-1)^{[\frac{n-2}{4}]} 2^{\frac{3n}{2}-2} - (-1)^{\frac{n}{2}} 2^{n-1} + 4U_{n-1}(2,5)\big).$$

Proof. Putting $m = 4$ in Theorem 2.4 we see that

$$\begin{aligned}
&\sum_{\substack{k=0\\4|k-2}}^{n} \binom{n}{k} (-1)^{\frac{k-2}{4}} 2^{n+\frac{3k}{2}} B_{n-k} \\
&= \frac{n}{2}\Big\{4^{n-1}\big((-1)^{[\frac{n-2}{4}]} 2^{\frac{n}{2}} - 2(-1)^{\frac{n}{2}}\big) + 2i\big((-2-4i)^{n-1} - (-2+4i)^{n-1}\big)\Big\}.
\end{aligned}$$

As

$$\begin{aligned}
&2i\big((-2-4i)^{n-1} - (-2+4i)^{n-1}\big) \\
&= -2^n i\big((1+2i)^{n-1} - (1-2i)^{n-1}\big) \\
&= 4 \cdot 2^n \cdot \frac{1}{\sqrt{2^2 - 4\times 5}}\Big\{\Big(\frac{2+\sqrt{2^2-4\times 5}}{2}\Big)^{n-1} - \Big(\frac{2-\sqrt{2^2-4\times 5}}{2}\Big)^{n-1}\Big\} \\
&= 2^{n+2} U_{n-1}(2,5),
\end{aligned}$$

the result follows from the above.

Similarly, putting $m = 3$ in Theorem 2.4 we deduce the following result.

Corollary 2.7. *For $n = 2, 4, 6, \ldots$ we have*

$$\sum_{\substack{k=0\\4|k-2}}^{n} \binom{n}{k} (-1)^{\frac{k-2}{4}} 2^{n-\frac{k}{2}} \cdot 3^k B_{n-k} = \frac{n}{2}\big((-1)^{[\frac{n-2}{4}]} 2^{\frac{n}{2}} \cdot 3^{n-1} + 12U_{n-1}(2,10)\big).$$

Theorem 2.5. *For $n \in \mathbb{N}$ we have*

$$\begin{aligned}
&\sum_{\substack{k=1\\4|k}}^{n} \binom{n}{k} B_{n-k}(x)((-1)^{\frac{k}{4}} 2^{\frac{k}{2}-1} - 1) \\
&= \frac{n}{8}\Big\{2(x-1)^{n-1} - 2x^{n-1} + (x+i)^{n-1} + (x-i)^{n-1} \\
&\qquad - (x-1+i)^{n-1} - (x-1-i)^{n-1}\Big\}.
\end{aligned}$$

Proof. From Lemma 2.1 we have

$$\begin{aligned}&\sum_{k=0}^{n-1}\binom{n}{k}B_k(x)((1\pm i)^{n-k}-(\pm i)^{n-k}-(\pm i)^{n-k}+(\pm i-1)^{n-k})\\&=n(x\pm i)^{n-1}-n(x\pm i-1)^{n-1}.\end{aligned}$$

Thus,

$$\begin{aligned}&2\sum_{\substack{k=0\\2|n-k}}^{n-1}\binom{n}{k}B_k(x)((i+1)^{n-k}-2i^{n-k}+(i-1)^{n-k})\\&=\sum_{k=0}^{n-1}\binom{n}{k}B_k(x)(1+(-1)^{n-k})((i+1)^{n-k}-2i^{n-k}+(i-1)^{n-k})\\&=\sum_{k=0}^{n-1}\binom{n}{k}B_k(x)((i+1)^{n-k}-2i^{n-k}+(i-1)^{n-k}\\&\qquad+(-i+1)^{n-k}-2(-i)^{n-k}+(-i-1)^{n-k})\\&=n(x+i)^{n-1}-n(x+i-1)^{n-1}+n(x-i)^{n-1}-n(x-i-1)^{n-1}.\end{aligned}$$

As

$$\begin{aligned}(i+1)^{2m}-2i^{2m}+(i-1)^{2m}&=(2i)^m-2(-1)^m+(-2i)^m\\&=\begin{cases}2((-1)^{\frac{m}{2}}2^m-1) & \text{if } 2\mid m,\\ 2 & \text{if } 2\nmid m,\end{cases}\end{aligned}\tag{2.3}$$

from the above we deduce

$$\begin{aligned}&4\sum_{\substack{k=0\\4|n-k}}^{n-1}\binom{n}{k}B_k(x)\big((-1)^{\frac{n-k}{4}}2^{\frac{n-k}{2}}-1\big)+4\sum_{\substack{k=0\\4|n-k-2}}^{n-1}\binom{n}{k}B_k(x)\\&=n\big\{(x+i)^{n-1}+(x-i)^{n-1}-(x-1+i)^{n-1}-(x-1-i)^{n-1}\big\}.\end{aligned}$$

Using Lemma 2.1 we see that

$$\begin{aligned}&4\sum_{\substack{k=0\\4|n-k}}^{n-1}\binom{n}{k}B_k(x)+4\sum_{\substack{k=0\\4|n-k-2}}^{n-1}\binom{n}{k}B_k(x)\\&=4\sum_{\substack{k=0\\2|n-k}}^{n-1}\binom{n}{k}B_k(x)=2\sum_{k=0}^{n-1}\binom{n}{k}B_k(x)(1^{n-k}-0^{n-k}-0^{n-k}+(-1)^{n-k})\\&=2nx^{n-1}-2n(x-1)^{n-1}.\end{aligned}$$

Therefore

$$4\sum_{\substack{k=0\\4|n-k}}^{n-1}\binom{n}{k}B_k(x)\big((-1)^{\frac{n-k}{4}}2^{\frac{n-k}{2}}-2\big)+2nx^{n-1}-2n(x-1)^{n-1}$$
$$=n\big\{(x+i)^{n-1}+(x-i)^{n-1}-(x-1+i)^{n-1}-(x-1-i)^{n-1}\big\}.$$

To see the result, we note that

$$\sum_{\substack{k=1\\4|k}}^{n}\binom{n}{k}B_{n-k}(x)\big((-1)^{\frac{k}{4}}2^{\frac{k}{2}}-2\big)=\sum_{\substack{k=0\\4|n-k}}^{n-1}\binom{n}{k}B_k(x)\big((-1)^{\frac{n-k}{4}}2^{\frac{n-k}{2}}-2\big).$$

Corollary 2.8. *For $n=2,4,6,\dots$ we have*

$$\sum_{\substack{k=1\\4|k}}^{n}\binom{n}{k}((-4)^{\frac{k}{4}}-2)B_{n-k}=\frac{n}{2}\big((-1)^{[\frac{n}{4}]}2^{\frac{n}{2}-1}-1\big).$$

Proof. Note that $2\nmid n-1$ and

$$(-1+i)^{n-1}+(-1-i)^{n-1}=(-1+i)(-2i)^{\frac{n-2}{2}}+(-1-i)(2i)^{\frac{n-2}{2}}=-(-1)^{[\frac{n}{4}]}2^{\frac{n}{2}}.$$

Now putting $x=0$ in Theorem 2.5 we deduce the result.

3. Recurrence formulas with gaps for Euler polynomials.

Lemma 3.1. *For any nonnegative integer n we have*

$$\sum_{k=0}^{n}\binom{n}{k}E_k(x)\big(z^{n-k}+(1+z)^{n-k}\big)=2(x+z)^n.$$

Proof. It is well known that

$$E_n(x+y)=\sum_{k=0}^{n}\binom{n}{k}E_n(x)y^{n-k}\quad\text{and}\quad E_n(x)+E_n(x+1)=2x^n.$$

Thus,

$$\sum_{k=0}^{n}\binom{n}{k}E_k(x)\big(z^{n-k}+(1+z)^{n-k}\big)=E_n(x+z)+E_n(x+z+1)=2(x+z)^n.$$

This is the result.

Theorem 3.1. *For $n \in \mathbb{N}$ we have*

$$\sum_{\substack{k=0\\4|k}}^{n}\binom{n}{k}(-1)^{\frac{k}{4}}2^{-\frac{k}{2}}E_{n-k}(x)=\frac{1}{2}\Big\{\Big(x-\frac{1}{2}+\frac{1}{2}i\Big)^n+\Big(x-\frac{1}{2}-\frac{1}{2}i\Big)^n\Big\}.$$

Proof. Putting $z=\frac{1}{2}(\pm i-1)$ in Lemma 3.1 we obtain

$$\sum_{k=0}^{n}\binom{n}{k}E_k(x)\big(\frac{1}{2}\big)^{n-k}\big((\pm i-1)^{n-k}+(1\pm i)^{n-k}\big)=2\Big(x+\frac{1}{2}(\pm i-1)\Big)^n.$$

Thus

$$\begin{aligned}&\sum_{k=0}^{n}\binom{n}{k}E_k(x)\big(\frac{1}{2}\big)^{n-k}\\&\qquad\times\Big\{(i-1)^{n-k}+(1+i)^{n-k}+(-i-1)^{n-k}+(1-i)^{n-k}\Big\}\\&=2\Big(x-\frac{1}{2}+\frac{1}{2}i\Big)^n+2\Big(x-\frac{1}{2}-\frac{1}{2}i\Big)^n.\end{aligned}$$

Clearly

$$\begin{aligned}&(i-1)^{n-k}+(1+i)^{n-k}+(-i-1)^{n-k}+(1-i)^{n-k}\\&=\big(1+(-1)^{n-k}\big)\big((i-1)^{n-k}+(1+i)^{n-k}\big)\\&=\big(1+(-1)^{n-k}\big)\big((-2i)^{\frac{n-k}{2}}+(2i)^{\frac{n-k}{2}}\big)\\&=\begin{cases}2^{\frac{n-k}{2}+2}(-1)^{\frac{n-k}{4}} & \text{if } 4\mid n-k,\\ 0 & \text{if } 4\mid n-k-2.\end{cases}\end{aligned}$$

So

$$\begin{aligned}&\sum_{\substack{k=0\\4|k}}^{n}\binom{n}{k}E_{n-k}(x)\big(\frac{1}{2}\big)^k(-1)^{\frac{k}{4}}2^{\frac{k}{2}+2}\\&=\sum_{\substack{k=0\\4|n-k}}^{n}\binom{n}{k}E_k(x)\big(\frac{1}{2}\big)^{n-k}(-1)^{\frac{n-k}{4}}2^{\frac{n-k}{2}+2}\\&=2\Big\{\Big(x-\frac{1}{2}+\frac{1}{2}i\Big)^n+\Big(x-\frac{1}{2}-\frac{1}{2}i\Big)^n\Big\}.\end{aligned}$$

This yields the result.

Corollary 3.1 ([L, (15)]). *For $n=2,4,6,\dots$ we have*

$$\sum_{\substack{k=0\\4|k}}^{n}\binom{n}{k}(-4)^{\frac{k}{4}}E_{n-k}=(-1)^{\frac{n}{2}}.$$

Proof. As $E_n=2^nE_n(\frac{1}{2})$, taking $x=\frac{1}{2}$ in Theorem 3.1 we deduce the result.

Corollary 3.2. *For $n = 2, 4, 6, \ldots$ we have*

$$\sum_{\substack{k=0\\4|k}}^{n-1}\binom{n}{k}(-1)^{\frac{k}{4}}2^{\frac{n-k}{2}}(2^{n-k}-1)B_{n-k} = (-1)^{[\frac{n}{4}]}\frac{n}{2}.$$

Proof. Taking $x = 0$ and replacing n with $n-1$ in Theorem 3.1 and then applying (1.4) we deduce the result.

We remark that Corollary 3.2 is equivalent to [S, Corollary 4.1].

Theorem 3.2. *For any nonnegative integer n we have*

$$\sum_{\substack{k=0\\4|k}}^{n}\binom{n}{k}\big((-1)^{\frac{k}{4}}2^{\frac{k}{2}-1}+1\big)E_{n-k}(x)$$
$$= \frac{1}{4}\big(2x^n + 2(x-1)^n + (x+i)^n + (x-i)^n + (x-1+i)^n + (x-1-i)^n\big).$$

Proof. From Lemma 3.1 we have

$$\sum_{k=0}^{n}\binom{n}{k}E_k(x)((1\pm i)^{n-k} + 2(\pm i)^{n-k} + (-1\pm i)^{n-k}) = 2(x\pm i)^n + 2(x-1\pm i)^n.$$

Thus,

$$\begin{aligned}
&2\sum_{\substack{k=0\\2|n-k}}^{n}\binom{n}{k}E_k(x)((i+1)^{n-k} + 2i^{n-k} + (i-1)^{n-k})\\
&= \sum_{k=0}^{n}\binom{n}{k}E_k(x)(1+(-1)^{n-k})((i+1)^{n-k} + 2i^{n-k} + (i-1)^{n-k})\\
&= \sum_{k=0}^{n}\binom{n}{k}E_k(x)((i+1)^{n-k} + 2i^{n-k} + (i-1)^{n-k}\\
&\qquad + (-i+1)^{n-k} + 2(-i)^{n-k} + (-i-1)^{n-k})\\
&= 2(x+i)^n + 2(x+i-1)^n + 2(x-i)^n + 2(x-i-1)^n.
\end{aligned}$$

As

$$\begin{aligned}
(i+1)^{2m} + 2i^{2m} + (i-1)^{2m} &= (2i)^m + 2(-1)^m + (-2i)^m\\
&= \begin{cases} 2((-1)^{\frac{m}{2}}2^m + 1) & \text{if } 2 \mid m,\\ -2 & \text{if } 2 \mid m-1,\end{cases}
\end{aligned}$$

from the above we deduce

$$\begin{aligned}
&4\sum_{\substack{k=0\\4|n-k}}^{n}\binom{n}{k}E_k(x)\big((-1)^{\frac{n-k}{4}}2^{\frac{n-k}{2}}+1\big) - 4\sum_{\substack{k=0\\4|n-k-2}}^{n}\binom{n}{k}E_k(x)\\
&= 2\big\{(x+i)^n + (x-1+i)^n + (x-i)^n + (x-1-i)^n\big\}.
\end{aligned}$$

Using Lemma 3.1 we see that

$$\begin{aligned}
&4\sum_{\substack{k=0\\4|n-k}}^{n}\binom{n}{k}E_k(x)+4\sum_{\substack{k=0\\4|n-k-2}}^{n}\binom{n}{k}E_k(x)\\
&=4\sum_{\substack{k=0\\2|n-k}}^{n}\binom{n}{k}E_k(x)=2\sum_{k=0}^{n}\binom{n}{k}E_k(x)(1^{n-k}+0^{n-k}+0^{n-k}+(-1)^{n-k})\\
&=2\big(E_n(x+1)+E_n(x)+E_n(x)+E_n(x-1)\big)\\
&=4x^n+4(x-1)^n.
\end{aligned}$$

Therefore

$$\begin{aligned}
&4\sum_{\substack{k=0\\4|n-k}}^{n}\binom{n}{k}E_k(x)\big((-1)^{\frac{n-k}{4}}2^{\frac{n-k}{2}}+2\big)\\
&=2\big((x+i)^n+(x+i-1)^n+(x-i)^n+(x-i-1)^n+2x^n+2(x-1)^n\big).
\end{aligned}$$

Replacing k with $n-k$ in the above formula we derive the result.

Theorem 3.3. *For any nonnegative integer n we have*

$$\begin{aligned}
&4E_n(x)+3\sum_{k=1}^{[n/6]}\binom{n}{6k}E_{n-6k}(x)\\
&=x^n+(x-1)^n+(-1)^nV_n(1-2x,x^2-x+1).
\end{aligned}$$

Proof. Taking $z=\omega,\omega^2$ in Lemma 3.1 we see that

$$\sum_{k=0}^{n}\binom{n}{k}E_k(x)\big(\omega^{n-k}+(1+\omega)^{n-k}\big)=2(x+\omega)^n$$

and

$$\sum_{k=0}^{n}\binom{n}{k}E_k(x)\big(\omega^{2(n-k)}+(1+\omega^2)^{n-k}\big)=2(x+\omega^2)^n.$$

Hence

$$\begin{aligned}
&\sum_{k=0}^{n}\binom{n}{k}E_k(x)\big(\omega^{n-k}+\omega^{2(n-k)}+(-\omega^2)^{n-k}+(-\omega)^{n-k}\big)\\
&=2(x+w)^n+2(x+\omega^2)^n.
\end{aligned}$$

Since

$$\begin{aligned}
&\omega^{n-k}+\omega^{2(n-k)}+(-\omega^2)^{n-k}+(-\omega)^{n-k}\\
&=(1+(-1)^{n-k})(\omega^{n-k}+\omega^{2(n-k)})=\begin{cases}0 & \text{if } 2\nmid n-k,\\ 4 & \text{if } 6\mid n-k,\\ 2(\omega+\omega^2)=-2 & \text{if } 6\mid n-k\pm 2,\end{cases}
\end{aligned}$$

we see that

$$\begin{aligned}
&6\sum_{\substack{k=0\\6|n-k}}^{n}\binom{n}{k}E_k(x)-2\sum_{\substack{k=0\\2|n-k}}^{n}\binom{n}{k}E_k(x)\\
&=\sum_{k=0}^{n}\binom{n}{k}E_k(x)(1+(-1)^{n-k})(\omega^{n-k}+\omega^{2(n-k)})\\
&=2(x+\omega)^n+2(x+\omega^2)^n.
\end{aligned}$$

That is,

$$6\sum_{\substack{k=0\\6|k}}^{n}\binom{n}{k}E_{n-k}(x)-2\sum_{\substack{k=0\\2|k}}^{n}\binom{n}{k}E_{n-k}(x)=2(x+\omega)^n+2(x+\omega^2)^n.$$

By Lemma 3.1 we have

$$\begin{aligned}
2E_n(x)+2\sum_{\substack{k=0\\2|n-k}}^{n}\binom{n}{k}E_k(x)&=\sum_{k=0}^{n}\binom{n}{k}E_k(x)(0^{n-k}+1^{n-k}+(-1)^{n-k}+0^{n-k})\\
&=2x^n+2(x-1)^n.
\end{aligned}$$

Thus,

$$\sum_{\substack{k=0\\2|k}}^{n}\binom{n}{k}E_{n-k}(x)=\sum_{\substack{k=0\\2|n-k}}^{n}\binom{n}{k}E_k(x)=x^n+(x-1)^n-E_n(x).$$

Hence,

$$\begin{aligned}
&6\sum_{k=0}^{[n/6]}\binom{n}{6k}E_{n-6k}(x)\\
&=2\sum_{k=0}^{[n/2]}\binom{n}{2k}E_{n-2k}(x)+2(x+\omega)^n+2(x+\omega^2)^n\\
&=2x^n+2(x-1)^n-2E_n(x)+2(x+\omega)^n+2(x+\omega^2)^n.
\end{aligned}$$

Therefore,

$$4E_n(x)=x^n+(x-1)^n+(x+\omega)^n+(x+\omega^2)^n-3\sum_{k=1}^{[n/6]}\binom{n}{6k}E_{n-6k}(x).$$

To see the result, we note that

$$\begin{aligned}
(-1)^n\big((x+\omega)^n+(x+\omega^2)^n\big)&=\Big(\frac{1-2x-\sqrt{-3}}{2}\Big)^n+\Big(\frac{1-2x+\sqrt{-3}}{2}\Big)^n\\
&=\Big(\frac{1-2x+\sqrt{(1-2x)^2-4(x^2-x+1)}}{2}\Big)^n\\
&\quad+\Big(\frac{1-2x-\sqrt{(1-2x)^2-4(x^2-x+1)}}{2}\Big)^n\\
&=V_n(1-2x,x^2-x+1).
\end{aligned}$$

Note that $E_n = 2^n E_n(\frac{1}{2})$. Putting $x = \frac{1}{2}$ in Theorem 3.3 we deduce (1.6).

References

[C] M. Chellali, *Accélération de calcul de nombres de Bernoulli*, J. Number Theory **28** (1988), 347-362.

[L] D.H. Lehmer, *Lacunary recurrence formulas for the numbers of Bernoulli and Euler*, Ann. Math. **36** (1935), 637-649.

[MOS] W. Magnus, F. Oberhettinger and R.P. Soni, *Formulas and Theorems for the Special Functions of Mathematical Physics* (3*rd Edition*), Springer-Verlag, New York, 1966, pp. 25-32.

[S] Z.H. Sun, *On the properties of Newton-Euler pairs*, J. Number Theory **114** (2005), 88-123.

[R] S. Ramanujan, *Some properties of Bernoulli's numbers*, J. Indian Math. Soc. **3** (1911), 219-234.

[W] H. C. Williams, *Édouard Lucas and Primality Testing, Canadian Mathematical Society Series of Monographs and Advanced Texts (Vol.22)*, Wiley, New York, 1998, pp. 74-92.